arXiv:1111.1709v1 [quant-ph] 7 Nov 2011

Dissipation-boosted entanglement in coupled harmonic oscillators

Chaitanya Joshi,[1, *] Michael J. W. Hall,[2, 3] Mats Jonson,[1, 4, 5] Patrik Öhberg,[1] and Erika Andersson[1]
[1] *SUPA, Department of Physics, Heriot-Watt University, Edinburgh, EH14 4AS, UK*
[2] *Theoretical Physics, RSPE, Australian National University, Canberra ACT 0200, Australia*
[3] *Centre for Quantum Dynamics, Griffith University, Brisbane, QLD 4111, Australia*
[4] *Department of Physics, University of Gothenburg, SE-412 96 Göteborg, Sweden*
[5] *Department of Physics, Division of Quantum Phases & Devices, Konkuk University, Seoul 143-701, Korea*

We investigate the effect of dissipation on indirectly coupled harmonic oscillators and show that the degree of entanglement of initially separable states, caused by squeezing, is enhanced by dissipation. This counterintuitive enhancement is generic, since it arises generally when squeezing is present, and squeezing is always present when the rotating-wave approximation is not made. The effect vanishes if all the oscillators are coupled to identical heat baths, suggesting that "heat flow" may be a necessary condition for dissipation-boosted quantum correlations in coupled harmonic oscillators.

PACS numbers: 03.67.Bg, 03.75.Gg, 85.85.+j

One of the strangest features of quantum mechanics, entanglement, is at the same time a crucial resource for quantum information processing [1–4]. It is well known that superposition states, including entangled states, are extremely sensitive to noise and dissipation. For example, decoherence induced by an environment tends to reduce quantum coherent superpositions to incoherent mixtures [5], an effect one typically wishes to minimize. Somewhat counterintuitively, however, decoherence can also be used to generate entanglement [6–8]. Schemes to that effect, typically involving atoms coupled to cavity fields, can be modeled as few-level quantum systems coupled to harmonic oscillators. In this Letter, in contrast, we are motivated by recent proposals to generate entangled states of micro- or nanomechanical oscillators [9–13]. We investigate the effect of dissipation on a set of pairwise linearly coupled harmonic oscillators, in a setting which does not involve any few-level systems.

Our aim is to investigate whether dissipation could be used to generate or enhance entanglement among coupled quantum harmonic oscillators initially prepared in "classical" separable Gaussian states, such as coherent or thermal states. Such a scheme could then be applied *e.g.* to micro- or nanomechanical oscillators coupled to cavity fields, and would be analogous to existing schemes involving quantum harmonic oscillators coupled to few-level systems. One easily realizes, however, that it is not possible to generate entanglement from classical initial states of quantum harmonic oscillators coupled by only passive, *i.e.,* nonsqueezing linear interactions. This is because the state of any number of passively coupled harmonic oscillator modes that are initially in a classical state will remain classical. This result still applies in the presence of decoherence and noise since this can be modeled via passive coupling to additional harmonic oscillator modes. Essentially, what one has, both in the presence and absence of dissipation, is equivalent to a linear optical network involving only beam splitters and phase shifters, which cannot generate any entanglement starting from classical states [14]. One therefore concludes that if all couplings are passive, the presence of few-level systems is crucial for dissipation to be able to enhance entanglement.

The above equivalence with linear optics, however, relies on making the rotating-wave approximation (RWA) [15], which can profoundly affect entanglement properties. For example, if one makes the RWA then the ground state of a chain of harmonic oscillators is separable, while if the RWA is not made, it is entangled [16]. More generally, if the RWA is not made for linearly coupled oscillators, then generic squeezing interactions remain and the coupling is no longer passive. In this case entanglement between the oscillators, initially prepared in "classical" states, may indeed arise. We have found that, remarkably, decoherence can then act to *increase* the degree of entanglement. This appears to be a stochastic resonance-like phenomenon, reminiscent of that found for a cavity QED setting [8], and it may be a general phenomenon. Interestingly, it seems that entanglement is enhanced only if the oscillators are coupled to baths with different temperatures and/or dissipation rates.

In this Letter we investigate in detail two indirectly coupled quantum harmonic oscillators, where the interaction between the two oscillators is mediated via linear couplings to a third middle oscillator. We will also remark briefly on the scenario in which the two oscillators are directly coupled to each other.

Denoting the vibron excitation creation (destruction) operators for the two end oscillators by $\hat{a}^\dagger(\hat{a})$ and $\hat{b}^\dagger(\hat{b})$, and for the mediating oscillator by $\hat{c}^\dagger(\hat{c})$, and assuming that all three oscillators are resonant, the Hamiltonian is given by

$$H = \hbar\omega(\hat{a}^\dagger\hat{a}+\hat{b}^\dagger\hat{b}+\hat{c}^\dagger\hat{c})+\hbar\kappa(\hat{a}+\hat{a}^\dagger+\hat{b}+\hat{b}^\dagger)(\hat{c}^\dagger+\hat{c}), \quad (1)$$

where κ is the coupling strength between end and middle oscillators. We assume that each oscillator is coupled to its own independent equilibrium heat bath. The time evolution of the system in the Born-Markov approximation is then described by the Lindblad-type master equation

$$\frac{\partial\rho}{\partial t} = -\frac{i}{\hbar}[H,\hat{\rho}] + \gamma\mathcal{L}_a\hat{\rho} + \gamma\mathcal{L}_b\hat{\rho} + \Gamma(\bar{n}+1)\mathcal{L}_c\hat{\rho} + \Gamma\bar{n}\mathcal{L}_{c^\dagger}\hat{\rho}, \tag{2}$$

where γ is the decay rate of the two end oscillators, coupled to zero-temperature heat baths, Γ is the decay rate of the mediating oscillator, which is coupled to a heat bath with average thermal occupancy $\bar{n}$, and $\mathcal{L}_x\hat{\rho} \equiv 2\hat{x}\hat{\rho}\hat{x}^\dagger - \hat{x}^\dagger\hat{x}\hat{\rho} - \hat{\rho}\hat{x}^\dagger\hat{x}$.

We shall solve the master equation (2) using the full Hamiltonian (1). For later reference we note that the RWA amounts to dropping terms with two annihilation or two creation operators, so that the interaction part of the Hamiltonian (1) reduces to

$$H_{\mathrm{RWA}} = \hbar\kappa(\hat{a}\hat{c}^\dagger + \hat{c}\hat{a}^\dagger + \hat{b}\hat{c}^\dagger + \hat{c}\hat{b}^\dagger). \tag{3}$$

The RWA is a good approximation if $\kappa \ll \omega$. It greatly simplifies analytical and numerical treatment, but fails to capture the quantum correlations that arise for classical states evolving according to (1).

Analyzing the entanglement dynamics for continuous variable systems is rather involved unless one only considers Gaussian states, which include thermal, squeezed and coherent states. This fits our purpose, since we consider classical initial states, and the Hamiltonian contains squeezing terms but no higher order terms. To solve the master equation (2) with the full Hamiltonian (1), we define a normal-ordered characteristic function [15] as $\chi(u,v,w) = \langle e^{u\hat{a}^\dagger}e^{-u^*\hat{a}}e^{v\hat{b}^\dagger}e^{-v^*\hat{b}}e^{w\hat{c}^\dagger}e^{-w^*\hat{c}}\rangle$. From the characteristic function one can obtain the expectation values of quantum mechanical observables, e.g. $\langle\hat{a}^{\dagger m}(t)\hat{b}^{\dagger n}(t)\rangle = (\frac{\partial}{\partial u})^m(\frac{\partial}{\partial v})^n\chi(u,v,w=0,t)|_{u,v=0}$.

If the oscillators are initially prepared in Gaussian states then it is consistent to make a Gaussian ansatz for the time-evolved characteristic function, $\chi(u,v,w,t) = \exp\left[-z^T\mathbf{A}(t)\,z + iz^Tb(t)\right]$. Here $\mathbf{A}(t)$ is a time dependent 4×4 symmetric matrix, $b(t)$ is a 4×1 time dependent vector and $z^T = (x_1,x_2,w_2,w_1)$ with $x_1 = (u+u^*+v+v^*)/4$, $x_2 = (u-u^*+v-v^*)/4i$ and $w = w_1 + iw_2$. The corresponding partial differential equation for $\chi(u,v,w,t)$ then becomes [15]

$$(\frac{\partial}{\partial t} + 2\kappa z^T\mathbf{N}z)\chi(u,v,w,t) = z^T\mathbf{M}\nabla\chi(u,v,w,t), \tag{4}$$

where $\nabla = (\frac{\partial}{\partial x_1},\frac{\partial}{\partial x_2},\frac{\partial}{\partial w_2},\frac{\partial}{\partial w_1})^T$ and

$$\mathbf{N} = \begin{pmatrix} \mathbf{0} & \mathbf{I} \\ \mathbf{I} & \bar{n}\Gamma/\kappa\mathbf{I} \end{pmatrix} \tag{5}$$

$$\mathbf{M} = \begin{pmatrix} -\gamma\mathbf{I}+\omega\mathbf{S} & \mathbf{U} \\ \mathbf{V} & -\Gamma\mathbf{I}-\omega\mathbf{S} \end{pmatrix}. \tag{6}$$

Here $\mathbf{0}$ is the 2×2 null matrix, $\mathbf{I}$ is a 2×2 identity matrix, and

$$\mathbf{S} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \mathbf{U} = \begin{pmatrix} 4\kappa & 0 \\ 0 & 0 \end{pmatrix}, \quad \mathbf{V} = \begin{pmatrix} 0 & 0 \\ 0 & 2\kappa \end{pmatrix}. \tag{7}$$

Recalling that $\mathbf{A}(t)$ is symmetric, we can write

$$\mathbf{A}(t) = \begin{pmatrix} \mathbf{P}(t) & \mathbf{Q}(t) \\ \mathbf{Q}(t)^T & \mathbf{R}(t) \end{pmatrix} \tag{8}$$

where $\mathbf{P}(t)$ and $\mathbf{R}(t)$ are 2×2 symmetric matrices. Using the above form of $\mathbf{A}(t)$, Eq. (4) reduces to the three coupled matrix differential equations

$$\begin{aligned} \dot{\mathbf{P}} &= \omega(\mathbf{SP}-\mathbf{PS}) - 2\gamma\mathbf{P} + \mathbf{UQ}^T + \mathbf{QU} \\ \dot{\mathbf{R}} &= \omega(\mathbf{RS}-\mathbf{SR}) - 2\Gamma\mathbf{R} + \mathbf{VQ} + \mathbf{Q}^T\mathbf{V} + 2\bar{n}\Gamma\mathbf{I} \\ \dot{\mathbf{Q}} &= \omega(\mathbf{SQ}+\mathbf{QS}) - (\Gamma+\gamma)\mathbf{Q} + \mathbf{PV} + \mathbf{UR} + 2\kappa\mathbf{I}, \end{aligned} \tag{9}$$

along with $\dot{b}(t)=\mathbf{M}b(t)$. This set of equations can be numerically solved for the time dependent matrices $\mathbf{P}(t)$, $\mathbf{R}(t)$ and $\mathbf{Q}(t)$, which together with the solution $b(t) = \exp(\mathbf{M}t)b(0)$ allows us to obtain the two-mode quantum characteristic function $\chi(u,v,w=0,t)$. For the steady state, which is the regime of current interest to us, an analytic expression for $\chi(u,v,w=0,t)$ can indeed be derived but it is rather lengthy and impenetrable.

Noting that an initial Gaussian state evolving according to Eq.(2) will always maintain its Gaussian character, allows us to fully characterize the quantum correlations between the two end oscillators in terms of their Wigner covariance matrix. In the steady state, the covariance matrix $\mathbf{V}$ is a 4×4 stationary symmetric matrix with $V_{i,j} = (\langle R_iR_j + R_jR_i\rangle)/2$ where $i,j\epsilon\{a,b\}$ and $R^T = (\hat{q}_a,\hat{p}_a,\hat{q}_b,\hat{p}_b)$. Here $\hat{q}_i$ and $\hat{p}_i$ are the position and momentum quadratures of the ith oscillator. A widely used entanglement measure is logarithmic negativity, which is an entanglement monotone and easy to compute. For a two-mode Gaussian continuous variable state characterized by the covariance matrix $\mathbf{V}$, the logarithmic negativity is computed as $\mathcal{N} = \mathrm{Max}[0, -\log(2\nu_-)]$ [3, 17, 18] where ν_- is the smallest of the symplectic eigenvalues of the covariance matrix given by $\nu_- = \sqrt{\sigma/2 - \sqrt{(\sigma^2 - 4\mathrm{Det}\mathbf{V})}/2}$. Here $\sigma = \mathrm{Det}\mathbf{A_1} + \mathrm{Det}\mathbf{B_1} - 2\mathrm{Det}\mathbf{C_1}$ and $\mathbf{V} = \begin{pmatrix} \mathbf{A_1} & \mathbf{C_1} \\ \mathbf{C_1^T} & \mathbf{B_1} \end{pmatrix}$, where $\mathbf{A_1}$ ($\mathbf{B_1}$) accounts for the local variances of mode $a(b)$, and $\mathbf{C_1}$ for the inter-mode correlations.

The steady state value of the logarithmic negativity for the two coupled oscillators was evaluated and is shown in Fig. 1. Remarkably, steady state entanglement between the two coupled oscillators is a non-monotonic function of the dissipation strengths Γ and γ. For a given dissipation strength γ, there is an optimal value of Γ for which entanglement between the two indirectly coupled oscillators is maximized. This counterintuitive effect is the main result of the paper and is a clear signature of the possibility of increasing the quantum correlations by purely incoherent means. Some insight into why a certain level of dissipation may enhance the entanglement can be obtained by transforming to the interaction picture, where the time evolution of operators



FIG. 1: (Color online) Logarithmic negativity $\mathcal{N}$, quantifying the entanglement between the two indirectly coupled harmonic oscillators in the steady state, plotted as a function of Γ and average thermal occupancy of the bath $\bar{n}$ where $\omega = 1$, $\kappa = \omega/3$ and $\gamma = \omega/10$.

$\hat{O}_I(t) = \exp(iHt/\hbar)\hat{O}\exp(-iHt/\hbar)$ is governed by the Hamiltonian (1). In this way the entangling effect of the Hamiltonian part of the evolution is separated from that of the dissipation. Without dissipation, the state $\hat{\rho}_I$ does not evolve and the entanglement induced by the Hamiltonian part of the evolution is manifested in the time-dependence of $\hat{a}_I(t), \hat{b}_I(t)$ and $\hat{c}_I(t)$. In the interaction picture, including dissipation, the state $\hat{\rho}_I(t)$ evolves according to

$$\frac{d}{dt}\hat{\rho}_I(t) = [\gamma\mathcal{L}_a^I + \gamma\mathcal{L}_b^I + \Gamma(\bar{n}+1)\mathcal{L}_c^I + \Gamma\bar{n}\mathcal{L}_{c^\dagger}^I]\hat{\rho}_I(t), \quad (10)$$

where $\mathcal{L}_a^I, \mathcal{L}_b^I, \mathcal{L}_c^I$ and $\mathcal{L}_{c^\dagger}^I$ are obtained by replacing the time independent $\hat{a}, \hat{b}, \hat{c}, \hat{c}^\dagger$ with $\hat{a}_I(t), \hat{b}_I(t), \hat{c}_I(t), \hat{c}_I^\dagger(t)$.

A full diagonalisation of the Hamiltonian in (1) will be given elsewhere [19]. The essential feature is that creation and annihilation operators are coupled during the time evolution, that is,

$$\begin{aligned}[\hat{a}_I(t), \hat{b}_I(t), \hat{c}_I(t), \hat{a}_I^\dagger(t), \hat{b}_I^\dagger(t), \hat{c}_I^\dagger(t)]^T = \\ \mathbf{F}\cdot[\hat{a}(0), \hat{b}(0), \hat{c}(0), \hat{a}^\dagger(0), \hat{b}^\dagger(0), \hat{c}^\dagger(0)]^T, \quad (11)\end{aligned}$$

where $\mathbf{F}$ is a time-dependent 6×6 matrix. As a result, the dissipation terms in the interaction picture will contain not just terms with one creation and one annihilation operator, but also terms with two creation or two annihilation operators. This may give rise to additional squeezing. The noise may therefore generate further entanglement over and above that generated by the Hamiltonian H in (1). This explanation also indicates that enhancement of entanglement through dissipation should be a general phenomenon for non-passive linear interactions, since whenever squeezing is present in a Hamiltonian, the dissipation terms in a similar interaction picture will in general contain additional squeezing terms.

On the other hand, no boosting of entanglement is expected for isotropic dissipation. This is the case for instance when all oscillators are coupled with identical decay rates to baths with the same nonzero temperature $\bar{n}$. The no boosting argument relies on the property

$$(\mathcal{L}_a + \mathcal{L}_b + \mathcal{L}_c)\rho = (\mathcal{L}_s + \mathcal{L}_f + \mathcal{L}_g)\rho, \quad (12)$$

where s, f, g refer to the independently evolving normal modes for the Hamiltonian (1) [19], with $\hat{s} = (\hat{a} - \hat{b})/\sqrt{2}$, $\hat{f} = (\hat{a}+\hat{b})/2 + \hat{c}/\sqrt{2}$ and $\hat{g} = (\hat{a}+\hat{b})/2 - \hat{c}/\sqrt{2}$. Thus, isotropic dissipation for the a, b, c modes corresponds to isotropic dissipation for the independently evolving normal modes s, f, g. Since such dissipation can only degrade any squeezing of the normal modes, where this squeezing is expected to correspond to enhancement of entanglement between the a and b modes, asymmetry between the reservoirs therefore seems necessary for dissipation to enhance quantum correlations between harmonic oscillators. This is intriguing, since it indicates a possible connection between enhancement of quantum correlations and heat flow.

When the RWA is made, the dissipation terms in the corresponding interaction picture only contain terms with one creation and one annihilation operator, in contrast to Eq. (11), and thus are not expected to create squeezing. This is consistent with the general argument that no entanglement can be generated if all couplings among system and environment modes are passive. An easy way to see this is to consider the Glauber-Sudarshan P-representation [15]. Any two-mode state $\hat{\rho}_{12}$ can be written in terms of a function $P(\alpha,\beta)$ as

$$\hat{\rho}_{12} = \int d^2\alpha d^2\beta P(\alpha,\beta)|\alpha\rangle_{11}\langle\alpha|\otimes|\beta\rangle_{22}\langle\beta|. \quad (13)$$

For a classical state, the P function is positive, and the state is separable. If the P function is negative, however, this is a signature of non-classical correlations in the system [21]. Coherent states are always transformed into other coherent states when the coupling is passive and linear [15]. For example, using a 50/50 beam splitter, $|\alpha\rangle_1\otimes|\beta\rangle_2$ may be transformed into $|(\alpha+\beta)/\sqrt{2}\rangle_1\otimes|(\alpha-\beta)/\sqrt{2}\rangle_2$. It therefore follows that under such couplings an initially positive P function will remain positive, meaning that a classical state will remain classical and separable [14]. This is true also when dissipation is included, since dissipation can be modeled via passive linear interactions with a reservoir [22].

Within the RWA an exact analytical solution for the master equation is also possible [19]. For example, if the two end oscillators with $\gamma = 0$ and $\bar{n} = 0$ are initially prepared in thermal states with mean occupancy $\bar{n}'$, the steady state is a thermal mixture of "dark states",

$$\hat{\rho} = \sum_{n=0}^{\infty}\frac{\bar{n}'^n}{(\bar{n}'+1)^{n+1}}|\Psi_n^-\rangle\langle\Psi_n^-|\otimes|0_c\rangle\langle 0_c|, \quad (14)$$

where $|\Psi_n^-\rangle = (\hat{s}^\dagger)^n|0\rangle_a|0\rangle_b/(\sqrt{n!})$ and $\hat{s} = (\hat{a}-\hat{b})/\sqrt{2}$. The dark states $|\Psi_n^-\rangle$ are so called since they are not affected by dissipation in mode $\hat{c}$. Each dark state individually is entangled, but interestingly, the mixture in Eq. (14) is nevertheless separable, which can be confirmed by checking that its P function is nonnegative.

We note that the RWA Hamiltonian (3) looks very similar to the one studied in [8], where it is found that noise does enhance steady-state entanglement even if the RWA is made. The main difference is that in [8], two cavity modes were interacting with a noise-driven two-level atom, leading to entanglement between the two cavity modes. In our case the atom is replaced by a third oscillator, and as we have argued above, an initially classical state of any number of passively coupled harmonic oscillator modes will always remain separable if the RWA is made. The presence of few-level quantum systems is therefore necessary if entanglement is to be created with only passive linear couplings, whether dissipation is present or not.

It is interesting to note that nonclassical correlations beyond entanglement, as quantified by Gaussian quantum discord [20], show similar behavior as entanglement. Details of this will be provided elsewhere [19], but the main result shows that these quantum correlations survive in the steady state.

Our results may be relevant, *e.g.*, for a chain of coupled nano- or micromechanical oscillators [9], for two nanocantilevers indirectly coupled to an ultracold Bose-Einstein condensate [13], or for a chain of trapped ions in a Paul trap [23]. Direct Coulomb interaction between three such ions can provide an interaction of the form in Eq. (1). Our study shows that dissipation can play a constructive role in enhancing quantum correlations in continuous variable systems, and suggests a general method for checking if this might be the case. We have also studied the dynamics of only two directly coupled harmonic oscillators, both of which are asymmetrically coupled to their independent reservoirs. This physical scenario is not exactly the same as discussed before. This is because the ground state of a chain of linearly coupled harmonic oscillators can only have nearest-neighbor entanglement [9]. In this case, too, we have confirmed the constructive role of local dissipation in maximizing quantum entanglement of coupled harmonic oscillators in the steady state, although the effect is not as unambiguous as with the two indirectly coupled oscillators described in this Letter.

In conclusion, we have studied the dynamics of two indirectly coupled harmonic oscillators initially prepared in classical Gaussian states, with interactions mediated via a third oscillator. If the rotating-wave approximation is not made, then quantum correlations are induced between the two end oscillators. Strikingly, the degree of quantum correlations is maximized for some optimal value of the dissipation rate Γ. The connection between heat flow and the enhancement of quantum correlations merits further investigation.

It is a pleasure to acknowledge fruitful discussions with J. Anders, S. M. Barnett, V. Kendon, G. J. Milburn and D. Vitali. C. J. acknowledges support from the ORS scheme, M. J. partial support from the Swedish VR and the Korean WCU program funded by MEST/NFR (Grant No. R31-2008-000-10057-0), and E.A. from EPSRC EP/G009821/1.

---

[*] Electronic address: cj53@hw.ac.uk

[1] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).

[2] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Rev. Mod. Phys.* **81**, 865 (1999).

[3] G. Adesso and F. Illuminati, *J. Phys. A* **40**, 7821 (2007).

[4] S. L. Braunstein and H. J. Kimble, *Phys. Rev. A* **61**, 042302 (2000); T. Tyc and B. C. Sanders, *Phys. Rev. A* **65**, 042310 (2002).

[5] W. H. Zurek, *Rev. Mod. Phys.* **75**, 715 (2003).

[6] M. B. Plenio, S. F. Huelga, A. Beige, and P. L. Knight, *Phys. Rev. A* **59**, 2468 (1999).

[7] A. Beige, S. Bose, D. Braun, S. F. Huelga, P. L. Knight, M. B. Plenio, and V. Vedral, *J. Mod. Opt.* **47**, 2583 (2000).

[8] M. B. Plenio and S. F. Huelga, *Phys. Rev. Lett.* **88,** 197901 (2002).

[9] J. Eisert, M. B. Plenio, S. Bose, and J. Hartley, *Phys. Rev. Lett.* **93,** 190402 (2004).

[10] S. Mancini, V. Giovannetti, D. Vitali, and P. Tombesi, *Phys. Rev. Lett.* **88**, 120401 (2002); S. Huang and G. S. Agarwal, *New J. Phys.* **11**, 103044 (2009).

[11] M. Ludwig, K. Hammerer, and F. Marquardt *Phys. Rev. A* **82**, 012333 (2010).

[12] F. Galve, L. A. Pachon, and D. Zueco, *Phys. Rev. Lett.* **105,** 180501 (2010).

[13] C. Joshi, A. Hutter, F. E. Zimmer, M. Jonson, E. Andersson and P. Öhberg, *Phys. Rev. A* **82**, 043846 (2010).

[14] M. S. Kim, W. Son, V. Buzek, and P. L. Knight, *Phys. Rev. A* **65,** 032323 (2002).

[15] S. M. Barnett and P. M. Radmore, *Methods in Theoretical Quantum Optics* (Oxford University Press, Oxford, 1997).

[16] J. Anders, *Phys. Rev. A* **77**, 062102 (2010).

[17] R. Simon, *Phys. Rev. Lett.* **84,** 2726 (2000).

[18] L.-M. Duan, G. Giedke, J. I. Cirac, and P. Zoller, *Phys. Rev. Lett.* **84,** 2722 (2000).

[19] C. Joshi, M. J. W. Hall, M. Jonson, P. Öhberg, and E. Andersson (In preparation).

[20] G. Adesso and A. Datta, *Phys. Rev. Lett.* **105,** 030501 (2010).

[21] L. Mandel, *Phys. Scr.* **T12**, 34 (1986).

[22] D. F. Walls, and G. J. Milburn, *Quantum Optics* (Springer Study Edition).

[23] K. R. Brown, C. Ospelkaus, Y. Colombe, A. C. Wilson, D. Leibfried, and D. J. Wineland, *Nature* **196,** 471 (2011).